

Evidence for infrared finite coupling in Sudakov resummation

Georges Grunberg

Centre de Physique Théorique de l'Ecole Polytechnique (CNRS UMR C7644),
91128 Palaiseau Cedex, France

(Dated: February 2, 2008)

New arguments are presented in favor of the infrared finite coupling approach to power corrections in the context of Sudakov resummation. The more regular infrared behavior of some peculiar combinations of Sudakov anomalous dimensions, free of Landau singularities at large N_f , is pointed out. A general conflict between the infrared finite coupling and infrared renormalon approaches to power corrections is explained, and a possible resolution is proposed, which makes use of the arbitrariness of the choice of constant terms in the Sudakov exponent. A simple ansatz for a ‘universal’ non-perturbative Sudakov effective coupling at large N_f emerges from these considerations. An alternative evidence for an infrared finite *perturbative* effective coupling in the Drell-Yan process at large N_f (albeit at odds with the infrared renormalon argument) is found within the framework of Sudakov resummation for eikonal cross sections of Laenen, Sterman and Vogelsang.

PACS numbers: 12.38.Cy, 11.15.Pg, 12.38.Aw

The notion of an infrared (IR) finite coupling, and the related concept of universality, to parametrize power corrections in QCD has attracted much attention for a long time [1]. In the present note I display further evidence in favor of this assumption in the more specific framework of Sudakov resummation, taking the examples of scaling violation in deep inelastic scattering (DIS) and in the Drell-Yan process (more details can be found in [2]). From the standard exponentiation formulas in Mellin N -space for the DIS structure function (I shall adopt in general the notations of [3]), one gets the large N relation

$$\frac{d \ln F_2(Q^2, N)}{d \ln Q^2} = 4C_F H(Q^2) + 4C_F \int_0^1 dz \frac{z^{N-1} - 1}{1 - z} A_S[(1 - z)Q^2] + \mathcal{O}(1/N), \quad (1)$$

where $H(Q^2)$ is given as a power series in $a_s(Q^2) \equiv \alpha_s(Q^2)/4\pi$ with N -independent coefficients, $4C_F A_S(k^2) = A(a_s(k^2)) + dB(a_s(k^2))/d \ln k^2$, and A (the universal ‘cusp’ anomalous dimension) and B are the standard Sudakov anomalous dimensions relevant to DIS, given as power series in a_s . $A_S(k^2)$ shall be referred to as the ‘Sudakov effective coupling’, a physical ‘effective charge’ [4] and we shall see (in the large N_f limit) that it has drastically different IR properties than the individual Sudakov anomalous dimensions it is composed of.

The Borel transform $B[A_S](u)$ of the Sudakov effective coupling has been computed [5] at large N_f . It is defined by $A_S(k^2) = \frac{1}{\beta_0} \int_0^\infty du \exp(-u \ln k^2/\Lambda^2) B[A_S](u)$, where $\beta_0 = \frac{11}{3}C_2(G) - \frac{2}{3}N_f$ is the one-loop coefficient of the beta function. One gets, with the ‘naïve non-abelization’ recipe: $B[A_S](u) = \exp(-du) \frac{\sin \pi u}{\pi u} \frac{1}{2} \left(\frac{1}{1-u} + \frac{1}{1-u/2} \right)$, where d is a scheme-dependent constant related to the renormalization of fermion loops: $d = -5/3$ in the $\overline{MS}$ scheme and $d = 0$ in the so called ‘V-scheme’. In the following I shall use the ‘V-scheme’ for simplicity. I

note that $B[A_S](u)$ is free of renormalon singularities, and that for $k = \Lambda$ (the Landau pole of the one-loop V-scheme coupling) $A_S(k^2 = \Lambda^2)$ is finite. This behavior is in striking contrast with that of the cusp anomalous dimension [6] which displays wild oscillations, resulting in a completely unphysical behavior around $k = \Lambda$. It is actually possible to get an analytic expression for $A_S(k^2)$, valid at all k^2 , and one finds that $A_S(k^2)$ approaches an *infinite* (and negative) IR fixed point for $k^2 \rightarrow 0$: $A_S(k^2) \simeq -\frac{1}{2\beta_0} \frac{\Lambda^4}{k^4} + \frac{1}{2\beta_0} \frac{\Lambda^2}{k^2}$. This is not by itself an unphysical behavior (the coupling is causal and there is no Landau singularity), except for the negative sign in the infrared, which cannot reproduce a vanishing Sudakov tail. The other trouble is in the too strongly divergent IR behavior, which gives a divergent contribution to the integral on the right-hand side of eq.(1). These facts make plausible the speculation that there exists a non-perturbative modification $\delta A_S(k^2)$ of the coupling at small scales [1] which might turn the infinite IR fixed point of perturbative origin into a genuinely non-perturbative, but softer (eventually finite) fixed point.

However, there is a potential clash between the IR finite coupling and the IR renormalons approaches to power corrections (very closely connected to the well-known issue [7],[6] of $1/Q$ corrections in Drell-Yan) which can be summarized as follows. Consider a typical ‘renormalon integral’ $R(Q^2, N) = \int_0^{Q^2} \frac{dk^2}{k^2} F(k^2/Q^2, N) A_S(k^2)$, where $F(k^2/Q^2, N)$ shall be referred to as the ‘Sudakov distribution function’, and introduce its (‘RS invariant’ [8]) Borel representation $R(Q^2, N) = \frac{1}{\beta_0} \int_0^\infty du \exp(-u \ln Q^2/\Lambda^2) B[R](u, N)$. The Borel transform of R is given by $B[R](u, N) = B[A_S](u) \times \tilde{F}(u, N)$, with $\tilde{F}(u, N) = \int_0^1 \frac{dx}{x} F(x, N) \exp(-u \ln x)$. The *factorized form* [8],[6] of the expression should be noted. Suppose now $B[A_S](u)$ has a zero at $u = u_0$, which is *not shared* by $B[R](u, N)$. Then $\tilde{F}(u, N)$ must have a pole at $u = u_0$. If $u_0 > 0$, this means an IR renormalon, and $F(x, N)$ contains an $\mathcal{O}(x^{u_0})$ contribution for $x \rightarrow 0$. If

this renormalon is not present in $B[R](u, N)$, the standard IR renormalon philosophy would conclude that no corresponding power correction is present. On the other hand, this power correction is still expected in the IR finite coupling approach, where $A_S(k^2)$ is assumed to have a finite IR fixed point and the low energy part of the renormalon integral $R(Q^2, N)$ is well-defined. Then the distribution function $F(k^2/Q^2, N)$ can be expanded in powers of k^2 for $k^2 \rightarrow 0$, yielding a non-vanishing power correction for each term [9] in its low energy expansion, parametrized by a low energy moment of the effective coupling $A_S(k^2)$. Even if $B[R](u, N)$ is singular, rather than non-vanishing and finite, at $u = u_0$, there is still a clash, since the singularity of $\tilde{F}(u, N)$ is necessarily stronger than that of $B[R](u, N)$ in presence of a zero of $B[A_S](u)$; thus the IR finite coupling approach will predict a *more enhanced* power correction than indicated by the renormalon argument. Two examples of this situation at large N_f are:

- i) *DIS*: there $B[A_S](u)$ vanishes for any integer u , $u \geq 3$.
- ii) *Drell-Yan*: in this case the Sudakov effective coupling which occurs in the Q^2 derivative of the Drell-Yan cross section is given by [10] $4C_F A_{S,DY}(k^2) = A(a_s(k^2)) + \frac{1}{2} dD_{DY}(a_s(k^2))/d \ln k^2$, and the corresponding Borel transform at $N_f = \infty$ is [6] (dropping an inessential $\exp(cu)$ factor which corresponds to a change of scale): $B[A_{S,DY}](u) = \frac{1}{\Gamma(1+u)} \frac{\pi^{1/2}}{\Gamma(1/2-u)}$, which vanishes for positive half-integer u . In both cases, the zeroes are absent from the corresponding Sudakov exponent, and the IR finite coupling approach will lead to the prediction of extra power corrections in the exponent (N^3/Q^6 , N^4/Q^8 , etc...in DIS, and N/Q , N^3/Q^3 , etc...in Drell-Yan). To make progress, we have to understand better the origin of zeroes in $B[A_S](u)$.

DIS case: putting $S(Q^2, N) = \int_0^1 dz z^{\frac{N-1}{1-z}-1} A_S[(1-z)Q^2]$, it is useful to write $S(Q^2, N)$ as a ‘renormalon integral’, where the Sudakov distribution function is given by $F(k^2/Q^2, N) = (1-k^2/Q^2)^{N-1} - 1$. Order by order the perturbative series of $S(Q^2, N)$ contain both $\mathcal{O}(N^0)$ ‘constant terms’ and terms which vanish as $N \rightarrow \infty$. A meaningful simplification is achieved by making use of the following important scaling property: defining $\epsilon = Nk^2/Q^2$, and $F(k^2/Q^2, N) \equiv G(\epsilon, N) = (1-\epsilon/N)^{N-1} - 1$, and taking the limit $N \rightarrow \infty$ with ϵ fixed one gets a *finite* result $G(\epsilon, N) \rightarrow G(\epsilon, \infty) \equiv G(\epsilon)$ with $G(\epsilon) = \exp(-\epsilon) - 1$. Let us now redefine the Sudakov exponent by using $G(\epsilon)$ as the new distribution function, i.e. $S(Q^2, N)$ is replaced by $S_{stan}(Q^2, N) = \int_0^{Q^2} \frac{dk^2}{k^2} G(Nk^2/Q^2) A_S(k^2)$ where the subscript ‘stan’ means ‘standard’. This step is legitimate since, order by order in perturbation theory, $S_{stan}(Q^2, N)$ and $S(Q^2, N)$ differ only [2] by terms which vanish as $N \rightarrow \infty$, and thus share the *same* $\ln N$ and constant terms. I next assume a similar ansatz for a modified Sudakov exponent

$$S_{new}(Q^2, N) = \int_0^{Q^2} \frac{dk^2}{k^2} G_{new}(Nk^2/Q^2) A_S^{new}(k^2), \quad (2)$$

and show that a *unique* solution for $G_{new}(Nk^2/Q^2)$ and $A_S^{new}(k^2)$ exists, under the condition that S_{new} reproduces all divergent $\ln N$ terms, together with an (a priori *arbitrary*) given set of constant terms. The latter are usually a subset of the set of all constant terms on the right hand side of eq.(1), which are uniquely defined, but (arbitrarily) split between those included in the Sudakov exponent $S(Q^2, N)$, and those belonging to $H(Q^2)$. This statement can be checked order by order in perturbation theory. Here I give an all-order proof. I first introduce a simplification appropriate to the large N limit. The perturbative series of $S_{new}(Q^2, N)$ still contains $\mathcal{O}(1/N)$ terms at large N . It is useful to discard them, and find a Borel representation of the corresponding series $S_{new}^{as}(Q^2, N)$ which contains only the (same) $\ln^p N$ and $\mathcal{O}(N^0)$ terms, with all the $\mathcal{O}(1/N)$ terms dropped. One can show the looked for Borel transform is given by

$$B[S_{new}^{as}](u, N) = B[A_S^{new}](u) \times \left[\exp(u \ln N) \int_0^\infty \frac{d\epsilon}{\epsilon} G_{new}(\epsilon) \exp(-u \ln \epsilon) - \frac{G_{new}(\infty)}{u} \right] \quad (3)$$

where $G_{new}(\infty) = -1$ is a subtraction term, and is fixed, since it determines the leading logs. Eq.(3) displays the general form of the large N Borel transform:

$$B[S_{new}^{as}](u, N) = \exp(u \ln N) B(u) + \frac{C_{new}(u)}{u}. \quad (4)$$

It is clear that the N -dependent part $B(u)$ of the Borel transform is unambiguously determined, and cannot be changed without changing the N -dependent terms, i.e. the coefficients of all logs of N , which are fixed. On the other hand, the constant terms included into the Sudakov exponent are obtained by setting $N = 1$ in eq.(3) or (4), and can be changed only by modifying the function $C_{new}(u)$ (with $C_{new}(0) = 1$). Given $B(u)$ from an independent calculation, and given an (a priori *arbitrary*) function $C_{new}(u)$, eq.(3) and (4) determine in principle *both* $B[A_S^{new}](u) = C_{new}(u)$ and $G_{new}(\epsilon)$, which completes the proof.

Large N_f result: Specializing now to large N_f to compute $B(u)$ (and eventually $C_{new}(u)$), the result for the Borel transform of $d \ln F_2(Q^2, N)/d \ln Q^2|_{SDG}$ at *finite* N can be given in the ‘massive’, ‘single dressed gluon’ (SDG) dispersive Minkowskian formalism [11],[1] as

$$B[d \ln F_2(Q^2, N)/d \ln Q^2]_{SDG}(u, N) = 4C_F \frac{\sin \pi u}{\pi u} \times \int_0^\infty \frac{dx}{x} \ddot{\mathcal{F}}_{SDG}(x, N) \exp(-u \ln x), \quad (5)$$

where the ‘characteristic function’ $\mathcal{F}_{SDG}(x, N)$ has been computed in [1] ($x = \lambda^2/Q^2$ where λ is the ‘gluon mass’). I have checked that here too a similar scaling property holds at large N , namely, putting $\mathcal{G}_{SDG}(y, N) \equiv \mathcal{F}_{SDG}(x, N)$ with $y \equiv Nx = N\lambda^2/Q^2$, one gets for $N \rightarrow$

∞ at fixed y : $\ddot{G}_{SDG}(y, N) \rightarrow \ddot{G}_{SDG}(y, \infty) \equiv \ddot{G}_{SDG}(y)$, with

$$\ddot{G}_{SDG}(y) = -1 + \exp(-y) - \frac{1}{2}y \exp(-y) - \frac{1}{2}y \Gamma(0, y) + \frac{1}{2}y^2 \Gamma(0, y), \quad (6)$$

where $\Gamma(0, y)$ is the incomplete gamma function and implies $\int_0^\infty \frac{dy}{y} \ddot{G}_{SDG}(y) \exp(-u \ln y) = \Gamma(-u) \frac{1}{2} \left(\frac{1}{1-u} + \frac{1}{1-u/2} \right)$. Thus for $N \rightarrow \infty$

$$B[d \ln F_2(Q^2, N)/d \ln Q^2]_{SDG}^{as}(u, N) = 4C_F \frac{\sin \pi u}{\pi u} \times \left[\exp(u \ln N) \int_0^\infty \frac{dy}{y} \ddot{G}_{SDG}(y) \exp(-u \ln y) + \frac{\Gamma_{SDG}(u)}{u} \right]$$

where $\Gamma_{SDG}(u) = \left(\frac{\sin \pi u}{\sin \pi u} \right)^2 \frac{1}{(1-u)(1-u/2)}$ is again a subtraction function which takes into account *all* constant terms of the SDG result. The latter are indeed obtained by setting $N = 1$ in eq.(7). Thus, if one wants to select as input an *arbitrary* subset of constant terms to be included into a *new* asymptotic Sudakov exponent $S_{new}^{as}(Q^2, N)$, one should simply change the subtraction function $\Gamma_{SDG}(u)$, namely define a new input $B[d \ln F_2(Q^2, N)/d \ln Q^2]_{SDG}^{as, new}$ by eq.(7), with $\Gamma_{SDG}(u) \rightarrow \Gamma_{new}(u)$ where $\Gamma_{new}(u)$ (with $\Gamma_{new}(0) = 1$) takes into account the new set of constant terms. Having no $\mathcal{O}(1/N)$ terms, one can now identify $4C_F B[S_{new}^{as}](u, N)$ with $B[d \ln F_2(Q^2, N)/d \ln Q^2]_{SDG}^{as, new}(u, N)$, which yields the *two* master equations

$$B[A_S^{new}](u) \int_0^\infty \frac{d\epsilon}{\epsilon} G_{new}(\epsilon) \exp(-u \ln \epsilon) \equiv B(u) = \frac{\sin \pi u}{\pi u} \int_0^\infty \frac{dy}{y} \ddot{G}_{SDG}(y) \exp(-u \ln y), \quad (8)$$

and

$$B[A_S^{new}](u) \equiv C_{new}(u) = \left(\frac{\sin \pi u}{\pi u} \right) \Gamma_{new}(u). \quad (9)$$

Eq.(8) and (9) allow to determine *both* the Sudakov distribution function $G_{new}(\epsilon)$ and the Borel transform of the associated Sudakov effective coupling $B[A_S^{new}](u)$, for a given input set of constant terms which fix the subtraction function $\Gamma_{new}(u)$. It is interesting that $B[A_S^{new}](u)$ is given *entirely* by the subtraction term. Applying these results to the standard exponentiation formula, where $G(\epsilon)$ is known, and using eq.(8), one can in particular rederive the previously quoted result for $B[A_S](u)$. I note that the simplest possible ansatz, namely $\Gamma_{simple}(u) \equiv 1$, gives $B[A_S^{simple}](u) = \frac{\sin \pi u}{\pi u}$, hence

$$A_S^{simple}(k^2) = \frac{1}{\beta_0} \left[\frac{1}{2} - \frac{1}{\pi} \arctan(t/\pi) \right], \quad (10)$$

with $t = \ln k^2/\Lambda^2$. It turns out that $A_S^{simple}(k^2)$ is already causal [2] and IR finite (with $A_S^{simple}(0) = 1/\beta_0$) at the *perturbative* level, and no non-perturbative modification is a priori necessary. However, the first two zeroes at $u = 1$ and $u = 2$ lead to two leading *log-enhanced* $\mathcal{O}(\frac{N}{Q^2} \ln Q^2)$ and $\mathcal{O}(\frac{N^2}{Q^4} \ln Q^2)$ power corrections at large N in the IR finite coupling framework, as can be easily checked since we have in this case $G_{simple}(\epsilon) = \ddot{G}_{SDG}(\epsilon) \simeq \frac{1}{2}[\epsilon(\ln \epsilon + \gamma_E - 3) + \epsilon^2(-\ln \epsilon - \gamma_E + 1) + \mathcal{O}(\epsilon^3)]$. There is thus, as expected, a discrepancy with the IR renormalon expectation, which predicts only $\mathcal{O}(\frac{N}{Q^2})$ and $\mathcal{O}(\frac{N^2}{Q^4})$ power corrections.

Drell-Yan case: The analogue of eq.(1) for the scaling violation of the short distance Drell-Yan cross-section is $d \ln \sigma_{DY}(Q^2, N)/d \ln Q^2 = 4C_F [H_{DY}(Q^2) + S_{DY}(Q^2, N)] + \mathcal{O}(1/N)$ with $(7) S_{DY}(Q^2, N) = \int_0^1 dz \frac{2z^{N-1}-1}{1-z} A_{S, DY}[(1-z)^2 Q^2]$. $S_{DY}(Q^2, N)$ can again be written as a ‘renormalon integral’, with $F(k^2/Q^2, N) \rightarrow F_{DY}(k^2/Q^2, N) = (1 - k/Q)^{N-1} - 1$ and $A_S(k^2) \rightarrow A_{S, DY}(k^2)$. The Sudakov distribution function $F_{DY}(k^2/Q^2, N)$ involves *both* [7] even and odd powers of k at small k . Taking the scaling limit $N \rightarrow \infty$ with $\epsilon_{DY} = Nk/Q$ fixed one thus gets $G_{DY}(\epsilon_{DY}) = \exp(-\epsilon_{DY}) - 1$. Now the work of [12] for eikonal cross sections suggests to use instead (thus avoiding odd power terms) $S_{DY}^{new}(Q^2, N) = \int_0^{Q^2} \frac{dk^2}{k^2} G_{DY}^{new}(Nk/Q) A_{S, DY}^{new}(k^2)$, with a distribution function (I deal with the log-derivative of the Drell-Yan cross-section)

$$G_{DY}^{new}(Nk/Q) = 2 \frac{d}{d \ln Q^2} [K_0(2Nk/Q) + \ln(Nk/Q) + \gamma_E], \quad (11)$$

i.e. $G_{DY}^{new}(\epsilon_{DY}) = -[1 + x \frac{dK_0}{dx}(x = 2\epsilon_{DY})]$, where $K_0(x)$ is the modified Bessel function of the second kind. I note that $G_{DY}^{new}(\epsilon_{DY}) \rightarrow -1$ for $\epsilon_{DY} \rightarrow \infty$, consistently with the large ϵ_{DY} limit of $G_{DY}(\epsilon_{DY})$. At large N_f one gets, instead of eq.(8) (eq.(9) remains the same)

$$B[A_{S, DY}^{new}](u) \int_0^\infty 2 \frac{d\epsilon}{\epsilon} G_{DY}^{new}(\epsilon) \exp(-2u \ln \epsilon) = \frac{\sin \pi u}{\pi u} \int_0^\infty \frac{dy}{y} \ddot{G}_{SDG, DY}(y) \exp(-u \ln y), \quad (12)$$

where $y = N^2 \lambda^2/Q^2$. On the other hand, the large N_f calculation of [6] yields the N -dependent part of the large N Borel transform

$$\frac{\sin \pi u}{\pi u} \int_0^\infty \frac{dy}{y} \ddot{G}_{SDG, DY}(y) \exp(-u \ln y) = -\frac{1}{u} \frac{\Gamma(1-u)}{\Gamma(1+u)}, \quad (13)$$

whereas eq.(11) gives

$$\int_0^\infty 2 \frac{d\epsilon}{\epsilon} G_{DY}^{new}(\epsilon) \exp(-2u \ln \epsilon) = -u [\Gamma(-u)]^2. \quad (14)$$

From eq.(12) one thus derives the large N_f result

$$B[A_{S,DY}^{new}](u) = \frac{1}{\Gamma(1+u)\Gamma(1-u)} \equiv \frac{\sin \pi u}{\pi u}, \quad (15)$$

hence $A_{S,DY}^{new}(k^2) = A_S^{simple}(k^2)$. Thus in the framework of [12] the *simplest* IR finite *perturbative* coupling naturally emerges! The fact that the new Sudakov distribution function $G_{DY}^{new}(\epsilon)$ implies a new Sudakov effective coupling is of course one of the main point of the present paper. Again, there is a discrepancy [13] with the IR renormalon expectation: although $G_{DY}^{new}(\epsilon_{DY}) \simeq \epsilon_{DY}^2 (2 \ln \epsilon_{DY} + 2\gamma_E - 1) + \mathcal{O}(\epsilon_{DY}^4 \ln \epsilon_{DY})$ involves only *even* powers of ϵ_{DY} for $\epsilon_{DY} \rightarrow 0$, they are logarithmically enhanced. One thus gets $\mathcal{O}(\frac{N^{2p}}{Q^{2p}} \ln Q^2)$ power corrections (p integer) at large N , instead of the $\mathcal{O}(\frac{N^{2p}}{Q^{2p}})$ corrections expected from IR renormalons.

Large N_f ansatz for a ‘universal’ Sudakov coupling: the question arises whether it is possible to reconcile the IR renormalon and IR finite coupling approaches. Actually, I should first stress it is not yet clear whether the two approaches should be necessarily reconciled. For instance, in the DIS case, it could be that the operator product expansion (OPE) at large N is consistent with the existence of higher order power corrections (N^3/Q^6 , N^4/Q^8 ,...) *in the exponent*. If this turns out to be the case, the IR finite coupling approach would be consistent with the OPE (and at odds with the IR renormalon prediction) even within the standard [3] exponentiation framework. Similarly, the ansatz $A_S^{simple}(k^2)$ might be the correct one in the Drell-Yan case [12], although it also contradicts the IR renormalon expectation. It is interesting to note that eq.(9) indicates that zeroes in $B[A_S^{new}](u)$ arise from *two* distinct sources: either the ‘universal’ $\sin \pi u / \pi u$ factor (*simple* zeroes at integer u can come only from there), or the ‘arbitrary’ $\Gamma_{new}(u)$ subtraction term (zeroes at half-integer u can come only from there). The previous discussion suggests that zeroes coming from the ‘universal’ $\sin \pi u / \pi u$ factor need *not* be necessarily removed in the IR finite coupling approach, at the difference of the more ‘artificial’ zeroes (such as those occurring in the standard Drell-Yan case) coming from the subtraction term.

Notwithstanding the above remarks, I shall adopt in the following the attitude that the IR renormalon and IR finite coupling approaches to power corrections should *always* be made consistent with one another. For this purpose, one must remove *all* zeroes from $B[A_S^{new}](u)$. The *mathematically simplest* ‘universal’ ansatz suggested by eq.(9) is to choose $\Gamma_{new}(u) = \Gamma(1-u)$, which yields $B[A_S^{new}](u) = (\frac{\sin \pi u}{\pi u}) \Gamma(1-u) = \frac{1}{\Gamma(1+u)}$. It is interesting to compare this result with the one obtained by eliminating the half-integer zeroes from the standard Drell-Yan result. There the simplest procedure is to define $B[A_{S,DY}^{new}](u) = \frac{\Gamma(1/2-u)}{\pi^{1/2}} B[A_{S,DY}](u)$, which yields again the previous ansatz. Given $B[A_S^{new}](u)$, eq.(8) or (12) determine the Sudakov distribution function $G_{new}(\epsilon)$. In particular, assuming the above ansatz, one gets in the DIS case

$$\begin{aligned} G_{new}(\epsilon) &= -\frac{\epsilon(1+\epsilon)}{2} \quad 0 \leq \epsilon \leq 1 \\ &= -1 \quad \epsilon \geq 1, \end{aligned} \quad (16)$$

where $\epsilon = Nk^2/Q^2$. Thus all power corrections beyond the two leading ones are indeed absent from the Sudakov exponent, in agreement with the renormalon argument. In the Drell-Yan case, one gets instead

$$G_{DY}^{new}(\epsilon_{DY}) = \exp(-\epsilon_{DY}^2) - 1, \quad (17)$$

i.e. only even powers of $\epsilon_{DY} = Nk/Q$, with no logarithmic enhancement for $\epsilon_{DY} \rightarrow 0$. There is compelling numerical evidence that the corresponding ‘universal’ Sudakov effective coupling: $A_S^{new}(k^2) = \frac{1}{\beta_0} \int_0^\infty du \exp(-u \ln k^2/\Lambda^2) \cdot 1/\Gamma(1+u)$ blows up very fast (but remains positive) for $k^2 \rightarrow 0$: $A_S^{new}(k^2) \simeq \frac{1}{\beta_0} \exp\left(\frac{\Lambda^2}{k^2}\right)$. Assuming that this behavior is indeed correct, one can again speculate that there exists a non-perturbative modification $\delta A_S^{new}(k^2)$ of the coupling at small scales which will generate a non-perturbative but finite IR fixed point (a simple ansatz for $\delta A_S^{new}(k^2)$ has been given in [2]).

To conclude, I have shown that in a number of examples at large N_f the Sudakov effective couplings display remarkably smooth IR properties, with no Landau singularities (a rather unusual occurrence in resummed perturbation theory). Generalizing from the present findings, one can distinguish two main possibilities. Either the subtraction term $\Gamma_{new}(u)$ has some singularities at finite and positive values of u (which must be simple poles for the corresponding Sudakov coupling to be free of IR renormalons): this is the case in DIS and Drell-Yan with the standard choice of the Sudakov distribution function; or it has no $u > 0$ singularities, as the ‘simple’ coupling eq.(10), which emerges naturally in the formalism of [12] for the Drell-Yan cross-section. In the former case, the corresponding Sudakov effective coupling is presumably [14] causal, but strongly IR divergent, requiring a low scale non-perturbative modification to build an IR finite coupling. In the latter case, the Sudakov coupling is likely (provided $\Gamma_{new}(u)$ does not decrease faster than an exponential for $u \rightarrow \infty$) to be already IR finite (and causal) at the *perturbative* level. Even then, there may be extra non-perturbative effects, which could also take the form of a non-perturbative modification of the coupling at low scales. I consider these facts as hints strongly supporting the IR finite coupling hypothesis. The above ansatzes are in conflict with the IR renormalon expectations for power corrections. The simplest attempt to reconcile the two approaches leads to the proposal of a *universal* perturbative Sudakov coupling, which again requires a non-perturbative modification to achieve IR finiteness. The latter could introduce some amount of non-universality. The variety of

Sudakov distribution functions showing up in the exponentiation procedure is related to the freedom to select an arbitrary set of ‘constant terms’ in the Sudakov exponent. This freedom leaves open the question of the determination of the ‘correct’ Sudakov distribution function. There is no obvious way to solve this ambiguity. In particular, one can show [2] that the natural option of including *all* $\mathcal{O}(N^0)$ terms into the Sudakov exponent is excluded in the DIS case: at large N_f , there is no solution for the corresponding Sudakov distribution function, if one defines the Sudakov effective coupling by eq.(9) with $\Gamma_{new}(u) \rightarrow \Gamma_{SDG}(u)$ (moreover this effective coupling turns out to have renormalons). In the Drell-Yan case, the corresponding Sudakov distribution function does exist, but implies $\mathcal{O}(1/Q)$ power corrections (and renormalons in the associated effective coupling). With the option to incorporate *all* constant terms into the Sudakov exponent disfavored, there remains the alternative to absorb into the exponent a ‘minimum’ set of constant terms, realized [15] by the (universal) choice $\Gamma_{new}(u) = 1$ corresponding to the ‘simple’ coupling eq.(10). A variant of this ‘minimal’ option, inspired by the work of [16], would allow to correlate the choice of ‘left-over’ constant terms functions H_{new} and H_{DY}^{new} not included in the exponents S_{new} and S_{DY}^{new} to the absolute ratio $|\frac{\mathcal{F}(-Q^2)}{\mathcal{F}(Q^2)}|$ of time-like and space-like quark form factors: one could require (at least at large N_f) that

$$4C_F(H_{DY}^{new}(Q^2) - 2H_{new}(Q^2)) = \frac{d}{d \ln Q^2} \left[\ln \left| \frac{\mathcal{F}(-Q^2)}{\mathcal{F}(Q^2)} \right|^2 \right].$$

Thus, given a ‘natural’ choice of G_{DY}^{new} , hence of H_{DY}^{new} , the corresponding DIS functions H_{new} and G_{new} would be fixed. An important issue is that of *universality* [17] (here extended between space-like and time-like processes): given an ansatz for a universal Sudakov effective coupling at large N_f (e.g. eq.(10)), one should check (work in progress) whether it remains universal at the perturbative level at *finite* N_f , using the *same* Sudakov distribution functions (e.g. eq.(6) and (11)) found at $N_f = \infty$. Further insight could be afforded by a better understanding of OPE at large N in the DIS case. Whatever the correct choice of the Sudakov distribution function turns out to be, it remains for future phenomenological work to determine the corresponding form of the (possibly non-perturbative) Sudakov coupling at *finite* N_f for each process, and test for eventual deviations from universality. This step requires a parametrization of the IR part of the coupling, which can be viewed as an alternative to the shape function approach [18], but does not involve any explicit IR cut-off.

I thank M. Beneke, Yu.L. Dokshitzer, J-P. Lansberg, G. Marchesini, G.P. Salam and G. Sterman for useful discussions. I am indebted to S. Friot for the result quoted in [15]. I also wish to thank the referee for thought-stimulating comments.

-
- [1] Yu.L. Dokshitzer, G. Marchesini and B.R. Webber, *Nucl. Phys.* **B469** (1996) 93, and references therein; Yu.L. Dokshitzer and B.R. Webber, *Phys.Lett.* **B404** (1997) 321.
 - [2] G. Grunberg, hep-ph/0601140.
 - [3] S. Moch, J.A.M. Vermaseren and A. Vogt, *Nucl.Phys.* **B726** (2005) 317.
 - [4] G. Grunberg, *Phys. Rev.* **D29** (1984) 2315.
 - [5] E. Gardi and R.G. Roberts, *Nucl.Phys.* **B653** (2003) 227.
 - [6] M. Beneke and V.M. Braun, *Nucl.Phys.* **B454** (1995) 253.
 - [7] G.P. Korchemsky and G. Sterman, *Nucl.Phys.* **B437** (1995) 415.
 - [8] G. Grunberg, *Phys.Lett.* **B304** (1993) 183.
 - [9] I am assuming here, as well as in eq.(1), that the effective coupling $A_S(k^2)$ arising directly from Sudakov resummation is an *Euclidean* coupling (rather than the (integrated) time-like discontinuity of an Euclidean coupling [1], which might be justified only in the single dressed gluon approximation), so that *all* power behaved terms (whether analytic or not) in the low energy expansion of $F(k^2/Q^2, N)$ do contribute to power corrections within the IR finite coupling framework. Identifying power corrections only through non-analytic terms will also lead to a conflict with the IR renormalon expectation if $B[A_S](u)$ does not vanish at the renormalon position.
 - [10] The natural occurrence of this combination of anomalous dimensions has been also pointed out by G.Sterman (talk given at the FRIF Workshop on first-principles non-perturbative QCD of hadron jets (Paris, January 2006)).
 - [11] M. Beneke and V.M. Braun, *Phys.Lett.* **B348** (1995) 513;
 - P. Ball, M. Beneke and V.M. Braun, *Nucl. Phys.* **B452** (1995) 563.
 - [12] G. Sterman and W. Vogelsang, hep-ph/9910371; E. Laenen, G. Sterman and W. Vogelsang, hep-ph/0010183; *Phys. Rev.* **D63** (2001) 114018.
 - [13] Consistently with the general assumption [9] that $A_{S,DY}^{new}(k^2)$ is an Euclidean coupling, I am discarding here (a similar remark applies below eq.(10)) the alternative interpretation that $G_{DY}^{new}(\epsilon_{DY})$ should be viewed as a ‘characteristic function’ [1] where power corrections are related to the *discontinuities* of non-analytic terms at small ϵ_{DY} .
 - [14] An exception is the case $C_{new}(u) = B[A_S^{new}](u) = 1$, i.e. $\Gamma_{new}(u) = \frac{\pi u}{\sin \pi u}$, which corresponds to a one loop effective Sudakov coupling.
 - [15] The alternative ‘minimal’ ansatz $B[A_S^{new}](u) = 1$ fits less well within the IR finite coupling approach, since it yields [14] an effective coupling with a Landau pole. Moreover, the corresponding Sudakov distribution function does *not* exist in the DIS case, although it does in the Drell-Yan case, where one gets $G_{DY}^{new}(\epsilon_{DY}) = J_0(2\epsilon_{DY}) - 1$ (J_0 is the Bessel function).
 - [16] G. Sterman, *Nucl.Phys.* **B281** (1987) 310; T.O. Eynck, E. Laenen and L. Magnea, *JHEP* **0306** (2003) 057.
 - [17] C. Lee and G. Sterman, hep-ph/0603066, and references therein.
 - [18] G.P. Korchemsky, hep-ph/9806537; G.P. Korchemsky and G. Sterman, *Nucl.Phys.* **B555** (1999) 335.